



## King's Research Portal

DOI:

[10.1016/j.ebiom.2017.08.006](https://doi.org/10.1016/j.ebiom.2017.08.006)

*Document Version*

Peer reviewed version

[Link to publication record in King's Research Portal](#)

*Citation for published version (APA):*

Nkya, S., Mgaya, J., Urrio, F., Makubi, A., Thein, S. L., Menzel, S., Cox, S. E., Newton, C. R., Kirkham, F. J., Mmbando, B. P., & Makani, J. (2017). Fetal Hemoglobin is Associated with Peripheral Oxygen Saturation in Sick Cell Disease in Tanzania. *EBioMedicine*. <https://doi.org/10.1016/j.ebiom.2017.08.006>

### **Citing this paper**

Please note that where the full-text provided on King's Research Portal is the Author Accepted Manuscript or Post-Print version this may differ from the final Published version. If citing, it is advised that you check and use the publisher's definitive version for pagination, volume/issue, and date of publication details. And where the final published version is provided on the Research Portal, if citing you are again advised to check the publisher's website for any subsequent corrections.

### **General rights**

Copyright and moral rights for the publications made accessible in the Research Portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognize and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the Research Portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the Research Portal

### **Take down policy**

If you believe that this document breaches copyright please contact [librarypure@kcl.ac.uk](mailto:librarypure@kcl.ac.uk) providing details, and we will remove access to the work immediately and investigate your claim.

## Accepted Manuscript

Fetal Hemoglobin is Associated with Peripheral Oxygen Saturation in Sickle Cell Disease in Tanzania

Siana Nkya, Josephine Mgaya, Florence Urio, Abel Makubi, Swee Lay Thein, Stephan Menzel, Sharon E. Cox, Charles R. Newton, Fenella J. Kirkham, Bruno P. Mmbando, Julie Makani



PII: S2352-3964(17)30323-7  
DOI: doi: [10.1016/j.ebiom.2017.08.006](https://doi.org/10.1016/j.ebiom.2017.08.006)  
Reference: EBIOM 1153

To appear in: *EBioMedicine*

Received date: 23 June 2017  
Revised date: 31 July 2017  
Accepted date: 5 August 2017

Please cite this article as: Siana Nkya, Josephine Mgaya, Florence Urio, Abel Makubi, Swee Lay Thein, Stephan Menzel, Sharon E. Cox, Charles R. Newton, Fenella J. Kirkham, Bruno P. Mmbando, Julie Makani, Fetal Hemoglobin is Associated with Peripheral Oxygen Saturation in Sickle Cell Disease in Tanzania. The address for the corresponding author was captured as affiliation for all authors. Please check if appropriate. *Ebiom*(2017), doi: [10.1016/j.ebiom.2017.08.006](https://doi.org/10.1016/j.ebiom.2017.08.006)

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Fetal hemoglobin is associated with peripheral oxygen saturation in sickle cell disease in Tanzania

Siana Nkya<sup>1,2</sup>, Josephine Mgaya<sup>1</sup>, Florence Urio<sup>1</sup>, Abel Makubi<sup>1</sup>, Swee Lay Thein<sup>3,4</sup>, Stephan Menzel<sup>3</sup>, Sharon E Cox<sup>1,5</sup>, Charles R Newton<sup>1,6</sup>, Fenella J Kirkham<sup>7,8,9</sup>, Bruno P Mmbando<sup>1,10</sup>, Julie Makani<sup>1,6</sup>

<sup>1</sup>Sickle cell Programme, Muhimbili University of Health and Allied Sciences, Dar-es-Salaam, Tanzania

<sup>2</sup>Dar-es-Salaam University College of Education, Dar-es-Salaam, Tanzania

<sup>3</sup>King's College London, Molecular Haematology, Division of Cancer Studies, UK

<sup>4</sup>Sickle cell branch, National heart, Lung and Blood Institute, The National Institutes of Health, USA

<sup>5</sup>Graduate School of Tropical Medicine & Global Health, Nagasaki University, Nagasaki, Japan, Faculty of Epidemiology & Population Health, London School of Hygiene & Tropical Medicine, London, UK

<sup>6</sup>Nuffield Department of Medicine, University of Oxford, UK

<sup>7</sup>UCL Great Ormond Street Institute of Child Health, London, UK

<sup>8</sup>Clinical and Experimental Sciences, University of Southampton, UK

<sup>9</sup>Southampton Children's Hospital, UK

<sup>10</sup>National Institute for Medical Research, Tanga Centre, Tanga, Tanzania,

#### Corresponding author:

Siana Nkya  
Department of Haematology and Blood Transfusion,  
P. O. Box 65001,  
Dar-es-Salaam, Tanzania.  
Phone: +255784193349  
Email: snkyamtatiro@gmail.com

#### ABSTRACT

Fetal hemoglobin (HbF) and peripheral hemoglobin oxygen saturation (SpO<sub>2</sub>) both predict clinical severity in sickle cell disease (SCD), while reticulocytosis is associated with vasculopathy, but there are few data on mechanisms. HbF, SpO<sub>2</sub> and routine clinical and laboratory measures were available in a Tanzanian cohort of 1175 SCD individuals aged ≥5 years and the association with SpO<sub>2</sub> (as response variable transformed to a Poisson distribution) was assessed by negative binomial model with age and sex as covariates. Increase in HbF was associated with increased SpO<sub>2</sub> (rate ratio, RR=1.19; 95% confidence intervals [CI] 1.04, 1.37 per natural log unit of HbF; P=0.0004). In

univariable analysis, SpO<sub>2</sub> was inversely associated with age, reticulocyte count, and log (total bilirubin) and directly with pulse, SBP, hemoglobin, and log (HbF). In multivariable regression log(HbF) (RR 1.191; 95%CI 1.04, 1.37; p=0.013), pulse (RR 1.01; 95%CI 1.00, 1.01; p=0.026), SBP (RR 1.008; 95%CI 1.00, 1.02; p=0.014), and hemoglobin (1.120; 95%CI 1.05, 1.19; p=0.001) were positively and independently associated with SpO<sub>2</sub> while reticulocyte count (RR 0.985; 95%CI 0.97, 0.99; p=0.019) was independently inversely associated with SpO<sub>2</sub>. In SCD, improving SpO<sub>2</sub>, in part through cardiovascular compensation and associated with reduced reticulocytosis, may be a mechanism by which HbF reduces disease severity.

**KEY WORDS:** Fetal hemoglobin (HbF), sickle cell disease, hypoxia, oxygen saturation, reticulocytes

#### HIGHLIGHTS

- Fetal hemoglobin may moderate sickle cell disease through increased oxygen saturation
- Low oxygen saturation is associated with reticulocytosis which might moderate cerebral vasculopathy and stroke risk
- Higher pulse rate and systolic blood pressure in those with higher SpO<sub>2</sub> suggests cardiovascular compensation for low SpO<sub>2</sub>

#### RESEARCH IN CONTEXT

Fetal hemoglobin (HbF) is normally synthesized during intrauterine life and it starts to decline before birth being replaced by adult hemoglobin (HbA). However some individuals continue to synthesize HbF to adulthood and are relatively protected from severe sickle cell disease. The mechanism of HbF protection in SCD has not been entirely established. This study reports a positive association between HbF and oxygen saturation (SpO<sub>2</sub>). Higher SpO<sub>2</sub> is associated with decreased reticulocytes but increased pulse rate and systolic blood pressure, suggesting SpO<sub>2</sub> is maintained in part through cardiovascular compensation. Increasing HbF may reduce disease severity partly through increasing SpO<sub>2</sub>.

#### INTRODUCTION

Sickle Cell disease (SCD) remains the most common hemoglobinopathy worldwide. Clinically, there is great variability amongst individuals with SCD; predictors of severity include hemoglobin F levels, reticulocytosis and alpha globin gene number. A major factor is the wide variation in the innate ability to synthesize fetal hemoglobin (HbF) beyond early childhood. Individuals with high levels of HbF experience milder forms of the disease with lower morbidity and improved survival. HbF level of 10% and above is believed to reduce the risk of major organ failure such as stroke, while much higher levels (20% and above) may be required to prevent recurrent clinical events such as painful crises and pulmonary disorder (Meier et al., 2017). The mechanisms underlying the reduction in the severity of SCD in people with high HbF are not clear. Therefore, studying the associations of HbF and clinical phenotypes of SCD may provide insights into the underlying mechanisms.

Peripheral hemoglobin oxygen saturation ( $SpO_2$ ), measured non-invasively by pulse oximetry, is related to several disease complications. Lower  $SpO_2$  has been associated with anemia (Quinn and Ahmad, 2005), increase in reticulocytes (Quinn and Ahmad, 2005), hemolysis (Campbell et al., 2009) and increased episodes of acute chest syndrome (Rackoff et al., 1993) and appears to predict central nervous system complications (including stroke (Quinn and Sargent, 2008), higher transcranial doppler velocity (Quinn et al., 2009), number of days per year admitted for pain (Hargrave et al., 2003), tricuspid regurgitant jet velocity (TRV) (Minniti et al., 2009) and diastolic dysfunction (Johnson et al., 2010) in SCD.

Studies in fetuses, children with cyanotic heart disease and adults ascending to high altitude suggest that HbF synthesis increases in hypoxia (Bard et al., 1994). Furthermore, hemolysis may induce HbF synthesis further (DeSimone et al., 1978). There is limited information on the magnitude and direction of any association between HbF and  $SpO_2$  in patients with SCD not on hydroxyurea, but the available data suggest that, as for neonates (Shiao, 2005),  $SpO_2$  is higher in SCD patients with higher HbF (Homi et al., 1997), (Halphen et al., 2014, Cox et al., 2013). In this report, we describe the association between HbF and  $SpO_2$  in individuals with SCD enrolled in the Muhimbili National Hospital cohort in Tanzania.

## MATERIALS AND METHODS

### Study area and population

This was a cross sectional study conducted at Muhimbili National Hospital (MNH) in Dar-es-Salaam, Tanzania involving individuals with SCD recruited into the Muhimbili Sickle cohort between March 2004 and December 2013. Recruitment and enrolment of patients and diagnosis of SCD has been previously described (Makani et al., 2011). Informed consent was obtained for each patient upon enrolment. Individuals were identified at pediatric SCD or hematology clinics or during hospitalization and were screened for SCD. A diagnosis of sickle cell anemia (HbSS/HbS $\beta^0$ ) by alkaline hemoglobin electrophoresis (Helena, Sunderland, Tyne & Wear, UK) was confirmed by high performance liquid chromatography (HPLC) (Variant I analyzer, Bio-Rad, Hercules, CA, USA). Ethical approval was granted by the Muhimbili University Research and Publications Committee (MU/RP/AEC/VOLX1/33).

Individuals were selected into this study if they had HbF values measured at the age of five years or above, since this is the age at which HbF synthesis stabilises. Data were excluded if the patient was on hydroxyurea therapy.

### Clinical measures

Daytime SpO<sub>2</sub> was determined in clinic when the child was well using a pulse oximeter (Nellcor, Pleasanton, CA, USA). Other clinical information that was collected included pulse rate, systolic blood pressure (SBP), and diastolic blood pressure (DBP).

### Laboratory measures

Hemoglobin was measured by an automated blood cell analyser (ABX Pentra 60 Analyser, Horiba, Kyoto, Japan) and reticulocytes were counted using the new methylene blue staining method followed by microscopy. HbF measurements were done by HPLC (Variant I, Biorad, Hercules, CA, USA). Routine biochemical analysis included total bilirubin (Roche Cobas Mira, New York, USA or Abbott Architect, New York, USA).

## Statistical methods

The SpO<sub>2</sub> data were collected as counts, which ranged from 82-100%, and could not be transformed into a normal random distribution, and hence a Poisson random distribution was assumed after transformation. To convert the distribution of SpO<sub>2</sub> (Figure 1A) into a Poisson distribution, 100-SpO<sub>2</sub> transformation was performed, Figure 1B. The distributions of HbF and bilirubin were positively skewed and hence normalized by natural log transformation. The association of clinical and laboratory variables with SpO<sub>2</sub> (as response variable) was assessed by negative binomial model with age and sex included as covariates, presenting the results as rate ratio (RR) with 95% confidence intervals. A p-value <0.05 was considered statistically significant. Variables with significant associations with SpO<sub>2</sub> in the univariate analysis were included in the multivariable regression analysis.

## RESULTS

We investigated 1,175 individuals (52.1% female, median age 11.2 [IQR: 7.9-16.7] years) with both HbF and SpO<sub>2</sub> measurements. Median HbF was 4.4% [IQR: 2.4-7.2] and median SpO<sub>2</sub> was 98% [IQR: 96-99]. Table 1 shows the results for the univariable and multivariable analyses. The distribution of HbF in relation to SpO<sub>2</sub> is presented in Figure 2; which shows the increase in oxygen saturation by one unit is associated with increase in mean of log(HbF) by 0.031, (95%CI: 0.013 0.049), p=0.020) while other variables are held constant (Table 1). In univariable analysis, SpO<sub>2</sub> was directly associated with pulse rate, systolic blood pressure (SBP), hemoglobin, and log (HbF) and inversely with age, reticulocyte count, and log (total bilirubin) (Table 1). In multivariable regression, log (HbF), pulse rate, SBP, and hemoglobin were positively and independently associated with SpO<sub>2</sub>, while reticulocyte count was inversely and independently associated with SpO<sub>2</sub> (Table 1).

## DISCUSSION

This study reports the association of HbF with SpO<sub>2</sub> in individuals with SCD. The study involves one of the largest single-centre SCD population to date and the first study to be conducted in an African population where the environment is different. This population is composed of a Central African Republic (CAR) sickle haplotype of predominately  $\beta^S/\beta^S$  genotype, a genetic background associated with a more severe disease. In addition, the population under study is Hydroxyurea naïve and there are limited resources for interventions. These factors make it pertinent for the study to be conducted in this population in order to identify disease-modifying factors that may be amenable to

treatment. We have confirmed the association between HbF and SpO<sub>2</sub> (Homi et al., 1997). Both HbF and total hemoglobin (Hb) were associated with SpO<sub>2</sub> independently suggesting that there may be separate mechanisms by which increasing Hb and HbF improve SpO<sub>2</sub>. The association between HbF and SpO<sub>2</sub> should augment efforts to develop and evaluate interventions that increase both HbF and SpO<sub>2</sub>. Although the mechanism of this association is not known, the oxygen dissociation curve properties for HbS and HbF are different. Therefore, the decreased oxygen affinity of hemoglobin S (Ueda et al., 1979) may partially contribute to the low SpO<sub>2</sub> in SCD. On the other hand, higher levels of HbF, which has a higher affinity for oxygen, results in a higher SpO<sub>2</sub>, in neonates (Shiao, 2005). In addition, compounds, including an aromatic aldehyde agent, 5-hydroxymethyl-2-furfural (5-HMF, also known as Aes-103), has been found to increase oxygen affinity of sickle hemoglobin and as a result reducing hypoxia-induced sickling *in vitro* and protects sickle cell mice from the effects of hypoxia (Safo and Kato, 2014).

Pulse rate, SBP and hemoglobin were also positively associated with SpO<sub>2</sub> in multivariable analysis. The direct association of hemoglobin with SpO<sub>2</sub> has been described previously and hence confirmed in the population that we have studied. The direct association of pulse rate and SBP suggests that one of the mechanisms for maintaining an adequate SpO<sub>2</sub> is cardiovascular compensation. The HbF levels for this population are low compared to other populations with different sickle haplotypes; despite this, an association with SpO<sub>2</sub> was established and can be further examined when interventions to increase HbF are in place. Treatment trials could examine whether improving SpO<sub>2</sub>, with agents that increase HbF or other methods of improving oxygenation, reduce SBP, a risk factor for stroke in adults and children with SCA.

An association between increased hemolysis and low SpO<sub>2</sub> has been reported (Nouraie et al., 2013). In this study, we report an inverse association between SpO<sub>2</sub> and total bilirubin and reticulocyte count, which are markers of hemolysis, on univariate analysis, suggesting higher SpO<sub>2</sub> in the presence of lower hemolysis or conversely, increased hemolysis if SpO<sub>2</sub> is lower. The effect of therapies designed to improve SpO<sub>2</sub> or reduce hemolysis may determine whether the initiating mechanism for the association involves either low SpO<sub>2</sub> or hemolysis. However, total bilirubin may also reflect liver compromise and reticulocytosis may be related to the response of the bone marrow to non-hemolytic anemia. On multivariable analysis, the reticulocyte count was independently associated with SpO<sub>2</sub>, which suggests an effect on erythropoiesis which may or may not involve hemolysis. We could not investigate the relative importance of hemolysis further as we only had



total bilirubin for one fifth of the patients; future studies should include measurement of indirect bilirubin, as a more specific marker of hemolysis, in all patients.

This study reports the association of HbF with SpO<sub>2</sub> two variables with strong clinical significance in individuals with SCD. The underlying mechanism of this association and the optimal range for HbF, measures cardiac function such as blood pressure and pulse, and SpO<sub>2</sub> for good health in SCD need to be established. This information will aid in the development and improvement of HbF-augmenting agents. The findings from this study may be applied to other SCD populations that may be similar.

## ACKNOWLEDGEMENTS

The authors thank the patients and staff of Muhimbili National Hospital, Muhimbili University of Health and Allied Sciences (MUHAS), Tanzania, Hematology Outpatient Unit and staff of the Muhimbili Wellcome Programme.

## FUNDING SOURCES

This work was supported by Wellcome Trust (Grant no: 084538).

The sponsors of this study are non-profit organizations that support science in general. They had no role in gathering, analyzing, or interpreting the data.

Bruno Mmbando is supported by NIH through H3Africa (4U41HG006941-02, <http://h3africa.org/>).

Siana Nkya conducted this work in partial fulfilment of the requirement for a PhD at Muhimbili University of Health and Allied Sciences.

## CONFLICT OF INTEREST

The authors declare no competing financial or other interests.

## AUTHOR CONTRIBUTIONS

Contribution: J.Makani, F.J.K, B.P. M and S.N.M. designed the study. J.Mgaya, collected the data. B.P.M. performed the analysis. S.N.M, J.Makani, B.P.M, S.C., C.R.N. and F.J.K wrote the manuscript and all authors commented on the drafts of the manuscript.

## REFERENCES

- BARD, H., FOURON, J. C., GAGNON, C. & GAGNON, J. (1994) Hypoxemia and increased fetal hemoglobin synthesis. *J Pediatr*, 124, 941-3.
- CAMPBELL, A., MINNITI, C. P., NOURAIE, M., ARTETA, M., RANA, S., ONYEKWERE, O., SABLE, C., ENSING, G., DHAM, N., LUCHTMAN-JONES, L., KATO, G. J., GLADWIN, M. T., CASTRO, O. L. & GORDEUK, V. R. (2009) Prospective evaluation of haemoglobin oxygen saturation at rest and after exercise in paediatric sickle cell disease patients. *Br J Haematol*, 147, 352-9.
- COX, S. E., MAKANI, J., NEWTON, C. R., PRENTICE, A. M. & KIRKHAM, F. J. (2013) Hematological and Genetic Predictors of Daytime Hemoglobin Saturation in Tanzanian Children with and without Sickle Cell Anemia. *ISRN Hematol*, 2013, 472909.
- DESIMONE, J., BIEL, S. I. & HELLER, P. (1978) Stimulation of fetal hemoglobin synthesis in baboons by hemolysis and hypoxia. *Proc Natl Acad Sci U S A*, 75, 2937-40.

- HALPHEN, I., ELIE, C., BROUSSE, V., LE BOURGEOIS, M., ALLALI, S., BONNET, D. & DE MONTALEMBERT, M. (2014) Severe nocturnal and postexercise hypoxia in children and adolescents with sickle cell disease. *PLoS One*, 9, e97462.
- HARGRAVE, D. R., WADE, A., EVANS, J. P., HEWES, D. K. & KIRKHAM, F. J. (2003) Nocturnal oxygen saturation and painful sickle cell crises in children. *Blood*, 101, 846-8.
- HOMI, J., LEEVE, L., HIGGS, D., THOMAS, P. & SERJEANT, G. (1997) Pulse oximetry in a cohort study of sickle cell disease. *Clin Lab Haematol*, 19, 17-22.
- JOHNSON, M. C., KIRKHAM, F. J., REDLINE, S., ROSEN, C. L., YAN, Y., ROBERTS, I., GRUENWALD, J., MAREK, J. & DEBAUN, M. R. (2010) Left ventricular hypertrophy and diastolic dysfunction in children with sickle cell disease are related to asleep and waking oxygen desaturation. *Blood*, 116, 16-21.
- MAKANI, J., COX, S. E., SOKA, D., KOMBA, A. N., ORUO, J., MWAMTEMI, H., MAGESA, P., RWEZAULA, S., MEDA, E., MGAYA, J., LOWE, B., MUTURI, D., ROBERTS, D. J., WILLIAMS, T. N., PALLANGYO, K., KITUNDU, J., FEGAN, G., KIRKHAM, F. J., MARSH, K. & NEWTON, C. R. (2011) Mortality in sickle cell anemia in Africa: a prospective cohort study in Tanzania. *PLoS One*, 6, e14699.
- MEIER, E. R., FASANO, R. M. & LEVETT, P. R. (2017) A systematic review of the literature for severity predictors in children with sickle cell anemia. *Blood Cells Mol Dis*, 65, 86-94.
- MINNITI, C. P., SABLE, C., CAMPBELL, A., RANA, S., ENSING, G., DHAM, N., ONYEKWERE, O., NOURAI, M., KATO, G. J., GLADWIN, M. T., CASTRO, O. L. & GORDEUK, V. R. (2009) Elevated tricuspid regurgitant jet velocity in children and adolescents with sickle cell disease: association with hemolysis and hemoglobin oxygen desaturation. *Haematologica*, 94, 340-7.
- NOURAI, M., LEE, J. S., ZHANG, Y., KANIAS, T., ZHAO, X., XIONG, Z., ORISS, T. B., ZENG, Q., KATO, G. J., GIBBS, J. S., HILDESHEIM, M. E., SACHDEV, V., BARST, R. J., MACHADO, R. F., HASSELL, K. L., LITTLE, J. A., SCHRAUFNAGEL, D. E., KRISHNAMURTI, L., NOVELLI, E., GIRGIS, R. E., MORRIS, C. R., ROSENZWEIG, E. B., BADESCH, D. B., LANZKRON, S., CASTRO, O. L., GOLDSMITH, J. C., GORDEUK, V. R., GLADWIN, M. T., WALK, P. I. & PATIENTS (2013) The relationship between the severity of hemolysis, clinical manifestations and risk of death in 415 patients with sickle cell anemia in the US and Europe. *Haematologica*, 98, 464-72.
- QUINN, C. T. & AHMAD, N. (2005) Clinical correlates of steady-state oxyhaemoglobin desaturation in children who have sickle cell disease. *Br J Haematol*, 131, 129-34.
- QUINN, C. T. & SARGENT, J. W. (2008) Daytime steady-state haemoglobin desaturation is a risk factor for overt stroke in children with sickle cell anaemia. *Br J Haematol*, 140, 336-9.
- QUINN, C. T., VARISTE, J. & DOWLING, M. M. (2009) Haemoglobin oxygen saturation is a determinant of cerebral artery blood flow velocity in children with sickle cell anaemia. *Br J Haematol*, 145, 500-5.
- RACKOFF, W. R., KUNKEL, N., SILBER, J. H., ASAKURA, T. & OHENE-FREMPONG, K. (1993) Pulse oximetry and factors associated with hemoglobin oxygen desaturation in children with sickle cell disease. *Blood*, 81, 3422-7.
- SAFO, M. K. & KATO, G. J. (2014). Therapeutic strategies to alter the oxygen affinity of sickle hemoglobin. *Hematol Oncol Clin North Am*, 28, 217-31.
- SHIAO, S. Y. (2005) Effects of fetal hemoglobin on accurate measurements of oxygen saturation in neonates. *J Perinat Neonatal Nurs*, 19, 348-61.
- UEDA, Y., NAGEL, R. L. & BOOKCHIN, R. M. (1979) An increased Bohr effect in sickle cell anemia. *Blood*, 53, 472-80.

Table 1: Association between SpO<sub>2</sub>(%) and clinical and laboratory parameters in Sickle cell disease

Parameters	N	SpO <sub>2</sub> (%) Univariable			SpO <sub>2</sub> (%), multivariable, N= 632		
		RR	95%CI	P	RR	95%CI	P
Age (years)	1175	0.99	0.89, 0.99	<b>0.024</b>	0.99	0.98, 1.01	0.422
Sex	1175	0.93	0.83, 1.04	0.191	0.91	0.78, 1.07	0.265
Pulse rate (beats/minute)	1170	1.00	1.00, 1.01	<b>0.020</b>	<b>1.01</b>	<b>1.00, 1.01</b>	<b>0.026</b>
Systolic Blood Pressure (mmHg)	1167	1.01	1.00, 1.01	<b>0.009</b>	<b>1.008</b>	<b>1.00, 1.02</b>	<b>0.014</b>
Log (HbF [%])	1175	1.19	1.08, 1.31	<b>0.0004</b>	<b>1.191</b>	<b>1.04, 1.37</b>	<b>0.013</b>
Hemoglobin (g/dL)	1136	1.07	1.07, 1.12	<b>0.001</b>	<b>1.120</b>	<b>1.05, 1.19</b>	<b>&lt;0.001</b>
Reticulocyte count (*10 <sup>9</sup> /L)	661	0.99	0.98, 1.00	<b>0.043</b>	<b>0.985</b>	<b>0.97, 0.99</b>	<b>0.019</b>
Log(Bilirubin total [mg/dl])	228	0.65	0.54, 0.77	<b>&lt;0.001</b>			

### Figure Legends

**Figure 1: Distribution of SpO<sub>2</sub> before (A) and after (B) transformation.** Conversion of the distribution of SpO<sub>2</sub> (Figure 1A) into a Poisson distribution, 100-SpO<sub>2</sub> transformation was performed resulting in the distribution shown in Figure 1B.

**Figure 2: Distribution of mean log (HbF) levels by Oxygen saturation.** The dotted line show the linear fit of the mean levels, which shows positive association with the increase with the oxy sat. The increase in oxy sat by one unit is associated with increase in mean of log(HbF) by 0.031, (95%CI: 0.013 0.049), p=0.020)



